



TITLE:

Spaces of holomorphic maps between complex projective spaces and group actions (New Evolution of Transformation Group Theory)

AUTHOR(S):

Yamaguchi, Kohhei

CITATION:

Yamaguchi, Kohhei. Spaces of holomorphic maps between complex projective spaces and group actions (New Evolution of Transformation Group Theory). 数理解析研究所講究録 2005, 1449: 72-81

ISSUE DATE:

2005-09

URL:

<http://hdl.handle.net/2433/47712>

RIGHT:

Spaces of holomorphic maps between complex projective spaces and group actions

電気通信大学 山口耕平 (Kohhei Yamaguchi)
University of Electro-Communications

1 Introduction.

The main purpose of this note is to announce the recent results concerning the topology of spaces of maps between complex projective spaces and related group actions on them.

First recall several notations used in this note. Let $j : S^2 = \mathbb{CP}^1 \rightarrow \mathbb{CP}^m$ be the inclusion map given by $j([x : y]) = [x : y : 0 : \cdots : 0]$. If $1 \leq m \leq n$ and $f : \mathbb{CP}^m \rightarrow \mathbb{CP}^n$ is a continuous map, the homotopy class of $f \circ j \in \pi_2(\mathbb{CP}^n) = \mathbb{Z}$ is called *the degree* of f . Let $\text{Map}_d(\mathbb{CP}^m, \mathbb{CP}^n)$ be the space consisting of all continuous maps $f : \mathbb{CP}^m \rightarrow \mathbb{CP}^n$ of degree d , and $\text{Map}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) \subset \text{Map}_d(\mathbb{CP}^m, \mathbb{CP}^n)$ the subspace of all based continuous maps $f : \mathbb{CP}^m \rightarrow \mathbb{CP}^n$ of degree d . We also denote by $\text{Hol}_d(\mathbb{CP}^m, \mathbb{CP}^n) \subset \text{Map}_d(\mathbb{CP}^m, \mathbb{CP}^n)$ (resp. $\text{Hol}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) \subset \text{Map}_d^*(\mathbb{CP}^m, \mathbb{CP}^n)$) the corresponding subspace consisting of all holomorphic maps (resp. based holomorphic maps). If $m > n$, there is no holomorphic map $\mathbb{CP}^m \rightarrow \mathbb{CP}^n$ except constant maps, and if $f : \mathbb{CP}^m \rightarrow \mathbb{CP}^n$ is a non-constant holomorphic map, $\deg f = d \geq 1$. So we shall only consider the case $1 \leq m \leq n$ with $d \geq 1$.

The origin of our work derives from the work of G. Segal [17], in which he describes that the Atiyah-Jones type result (cf. [1]) holds for the inclusion map $\text{Hol}(\mathbb{CP}^1, \mathbb{CP}^n) \rightarrow \text{Map}(\mathbb{CP}^1, \mathbb{CP}^n)$. For understanding his result intuitively, recall the case $m = n = 1$.

Let $C_d^\infty(\mathbb{CP}^1, \mathbb{CP}^1)$ denote the space of all smooth maps $f : \mathbb{CP}^1 \rightarrow \mathbb{CP}^1$ of degree d and consider the energy functional $E : C_d^\infty(\mathbb{CP}^1, \mathbb{CP}^1) \rightarrow \mathbb{R}$ defined by

$$E(f) = \frac{1}{2} \int_{\mathbb{CP}^1} \|df\|^2 d\text{vol} \quad \text{for } f \in C_d^\infty(\mathbb{CP}^1, \mathbb{CP}^1).$$

The critical point of E is called a *harmonic map* and it is known that the space of all harmonic maps is just $\text{Hol}_d(\mathbb{CP}^1, \mathbb{CP}^1)$ in this case. If we believe that Morse theoretical principle would hold for this case, $\text{Hol}_d(\mathbb{CP}^1, \mathbb{CP}^1) \subset C_d^\infty \simeq \text{Map}_d(\mathbb{CP}^1, \mathbb{CP}^1)$ is a deformation retract. This is clearly false, but Segal showed that it was true if " $d \rightarrow \infty$ ". More precisely, he proved the following important result:

Theorem 1.1 (G. Segal, [17]). *The inclusion maps*

$$\begin{cases} i_d : \text{Hol}_d^*(\mathbb{CP}^1, \mathbb{CP}^n) \rightarrow \text{Map}_d^*(\mathbb{CP}^1, \mathbb{CP}^n) = \Omega_d^2 \mathbb{CP}^n \simeq \Omega^2 S^{2n+1} \\ j_d : \text{Hol}_d(\mathbb{CP}^1, \mathbb{CP}^n) \rightarrow \text{Map}_d(\mathbb{CP}^1, \mathbb{CP}^n) \end{cases}$$

are homotopy equivalences up to dimension $(2n - 1)d$. \square

Remark. A map $f : X \rightarrow Y$ is called a *homotopy equivalence* (resp. a *homology equivalence*) up to dimension D if the induced homomorphism $f_* : \pi_k(X) \rightarrow \pi_k(Y)$ (resp. $f_* : H_k(X, \mathbb{Z}) \rightarrow H_k(Y, \mathbb{Z})$) is bijective when $k < D$ and surjective when $k = D$. Similarly, a map $f : X \rightarrow Y$ is called a *homotopy equivalence* (resp. a *homology equivalence*) through dimension D if the induced homomorphism $f_* : \pi_k(X) \rightarrow \pi_k(Y)$ (resp. $f_* : H_k(X, \mathbb{Z}) \rightarrow H_k(Y, \mathbb{Z})$) is bijective whenever $k \leq D$.

In [17], Segal also expected that a similar Atiyah-Jones type result would hold for the inclusion $\text{Hol}(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \text{Map}(\mathbb{CP}^m, \mathbb{CP}^n)$ even if $2 \leq m \leq n$, and we would like to investigate this problem. Recently we obtain the following result:

Theorem 1.2 ([12], [22]). *If $2 \leq m \leq n$, the inclusion maps*

$$\begin{cases} i_d : \text{Hol}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \text{Map}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) \\ j_d : \text{Hol}_d(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \text{Map}_d(\mathbb{CP}^m, \mathbb{CP}^n) \end{cases}$$

are homotopy equivalences through dimension $D(d; m, n)$, where $[x]$ denotes the integer part of a number x and the number $D(d; m, n)$ is given by $D(d; m, n) = (2n - 2m + 1) \left(\left\lfloor \frac{d+1}{2} \right\rfloor + 1 \right) - 1$.

2 The case $d = 1$.

Because the spaces $\text{Hol}_d(\mathbb{CP}^m, \mathbb{CP}^n)$ and $\text{Map}_d(\mathbb{CP}^m, \mathbb{CP}^n)$ are easily understood when $d = 1$, in this section we review the case $d = 1$.

First, we recall the interesting result due to S. Sasao [16].

Define the map $s'_{m,n} : U_{n+1} \rightarrow \text{Map}_1(\mathbb{CP}^m, \mathbb{CP}^n)$ by the usual matrix multiplication

$$s'_{m,n}(A)([x_0 : \cdots : x_m]) = [x_0 : \cdots : x_m : 0 : \cdots : 0] \cdot A$$

for $([x_0 : \cdots : x_m], A) \in \mathbb{CP}^m \times U_{n+1}$.

If $\Delta_k \subset U_k$ denotes the center of U_k , because the subgroup $\Delta_{m+1} \times U_{n-m} \subset U_{n+1}$ is fixed by the map $s'_{m,n}$, it induces the map

$$s_{m,n} : \text{PW}_{n+1,m+1} \rightarrow \text{Map}_1(\mathbb{CP}^m, \mathbb{CP}^n),$$

where $\text{PW}_{n+1,m+1} = U_{n+1}/(\Delta_{m+1} \times U_{n-m})$ denotes the complex projective Stiefel manifold of orthogonal $(m+1)$ -frames in \mathbb{C}^{n+1} . Then Sasao showed the following interesting result.

Theorem 2.1 (S. Sasao, [16]). *If $1 \leq m \leq n$, the map $s_{m,n} : \text{PW}_{n+1,m+1} \rightarrow \text{Map}_1(\mathbb{CP}^m, \mathbb{CP}^n)$ is a homotopy equivalence up to dimension $(4n - 4m + 1)$.* \square

Similarly, define the map $s''_{m,n} : U_n \rightarrow \text{Map}_1^*(\mathbb{CP}^m, \mathbb{CP}^n)$ by the matrix multiplication

$$s''_{m,n}(A)([x_0 : \cdots : x_m]) = [x_0 : \cdots : x_m : 0 : \cdots : 0] \cdot \begin{bmatrix} 1 & \mathbf{0}_n \\ {}^t\mathbf{0}_n & A \end{bmatrix}$$

for $([x_0 : \cdots : x_m], A) \in \mathbb{CP}^m \times U_n$, where $\mathbf{0}_n = (0, \dots, 0) \in \mathbb{C}^n$. Since the subgroup $U_{n-m} \subset U_n$ is fixed under this map, it also induces the map

$$\bar{s}_{m,n} : W_{n,m} \rightarrow \text{Map}_1^*(\mathbb{CP}^m, \mathbb{CP}^n),$$

where $W_{n,m} = U_n/U_{n-m}$ denotes the complex Stiefel manifold of orthogonal m -frames in \mathbb{C}^n . Then we also obtain:

Theorem 2.2. *If $1 \leq m \leq n$, the map $\bar{s}_{m,n} : W_{n,m} \rightarrow \text{Map}_1^*(\mathbb{CP}^m, \mathbb{CP}^n)$ is a homotopy equivalence up to dimension $4n - 4m + 1$.*

Proof. An easy diagram chasing shows that there is a commutative diagram of fibrations

$$\begin{array}{ccccc}
 W_{n,m} & \longrightarrow & PW_{n+1,m+1} & \longrightarrow & \mathbb{CP}^n \\
 \bar{s}_{m,n} \downarrow & & s_{m,n} \downarrow & & \parallel \\
 \text{Map}_1^*(\mathbb{CP}^m, \mathbb{CP}^n) & \xrightarrow{\subset} & \text{Map}_1(\mathbb{CP}^m, \mathbb{CP}^n) & \xrightarrow{ev} & \mathbb{CP}^n
 \end{array}$$

Then the assertion follows from the Five Lemma and the homotopy exact sequences of fibrations. \square

Now we can easily understand the reason why the two spaces $PW_{n+1,m+1}$ and $W_{n,m}$ approximate $\text{Map}_1(\mathbb{CP}^m, \mathbb{CP}^n)$ and $\text{Map}_1^*(\mathbb{CP}^m, \mathbb{CP}^n)$ up to homotopy equivalence as above. This can be explained by the following result.

Theorem 2.3 ([10]). *Let $1 \leq m \leq n$ and $d \geq 1$ be integers.*

(1) *If $d = 1$, there are homotopy equivalences*

$$\begin{cases} \text{Hol}_1^*(\mathbb{CP}^m, \mathbb{CP}^n) \simeq W_{n,m} \\ \text{Hol}_1(\mathbb{CP}^m, \mathbb{CP}^n) \simeq PV_{n+1,m+1} \end{cases}$$

(2) *The inclusion maps*

$$\begin{cases} i_1 : \text{Hol}_1^*(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \text{Map}_1^*(\mathbb{CP}^m, \mathbb{CP}^n) \\ j_1 : \text{Hol}_1(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \text{Map}_1(\mathbb{CP}^m, \mathbb{CP}^n) \end{cases}$$

are homotopy equivalence up to dimension $4n - 4m + 1$.

Sketch proof. We shall only give the proof of the existence of the homotopy equivalence $\text{Hol}_1^*(\mathbb{CP}^m, \mathbb{CP}^n) \simeq W_{n,m}$.

We take $e_j = [1 : 0 : 0 : \cdots : 0] \in \mathbb{CP}^j$ as the basepoint of \mathbb{CP}^j . An element $f \in \text{Hol}_d^*(\mathbb{CP}^m, \mathbb{CP}^n)$ is a holomorphic map $\mathbb{CP}^m \rightarrow \mathbb{CP}^n$ satisfying the condition $f(e_m) = e_n$. Such a map can be identified with the $(n+1)$ -tuple $f = (f_0, f_1, \dots, f_n)$ of homogeneous polynomials of the degree d in $\mathbb{C}[z_0, \dots, z_m]$ such that,

- (i) f_0, f_1, \dots, f_n have no common root except $0_{m+1} \in \mathbb{C}^{m+1}$,
- (ii) the coefficient of $(z_0)^d$ in f_0 is 1, and these in the other polynomials are 0.

If $d = 1$, we can write

$$\begin{aligned} f = (f_0, f_1, \dots, f_n) &= (z_0 + \sum_{k=1}^m b_k z_k, \sum_{k=1}^m a_{k,1} z_k, \dots, \sum_{k=1}^m a_{k,n} z_k) \\ &= (z_0, z_1, \dots, z_m) \cdot \begin{bmatrix} 1 & 0_n \\ {}^t\mathbf{b} & A \end{bmatrix}, \end{aligned}$$

where $\mathbf{b} = (b_1, \dots, b_m) \in \mathbb{C}^m$, $A = (a_{i,j})$: $(m \times n)$ -matrix.

First, we note that the polynomials f_0, f_1, \dots, f_n have no common root except 0_{m+1} if and only if $\text{rank } A = m$. Next, we also remark that the space of all $(m \times n)$ matrices of rank m is homeomorphic to the homogenous space $\text{GL}_n(\mathbb{C})/\text{GL}_{n-m}(\mathbb{C})$. Via this identification, we define a map $\gamma : \text{Hol}_1^*(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \mathbb{C}^m \times (\text{GL}_n(\mathbb{C})/\text{GL}_{n-m}(\mathbb{C}))$ by

$$\gamma(f) = \gamma(f_0 : f_1 : \dots : f_n) = (\mathbf{b}, A).$$

Next define a map $\beta' : \mathbb{C}^m \times \text{GL}_n(\mathbb{C}) \rightarrow \text{Hol}_1^*(\mathbb{CP}^m, \mathbb{CP}^n)$ by matrix multiplication

$$\beta'(\mathbf{b}, B)([x_0 : x_1 : \dots : x_m]) = [x_0 : x_1 : \dots : x_m : 0 : \dots : 0] \cdot \begin{bmatrix} 1 & 0_n \\ {}^t\tilde{\mathbf{b}} & B \end{bmatrix}$$

for $([x_0 : x_1 : \dots : x_m], \mathbf{b}, B) \in \mathbb{CP}^m \times \mathbb{C}^m \times \text{GL}_n(\mathbb{C})$, where $\tilde{\mathbf{b}} = (\mathbf{b}, 0_{n-m}) \in \mathbb{C}^n$. Since β' maps the subspace $\{0_m\} \times \text{GL}_{n-m}(\mathbb{C}) \subset \mathbb{C}^m \times \text{GL}_n(\mathbb{C})$ to the basepoint, it induces a map

$$\beta : \mathbb{C}^m \times (\text{GL}_n(\mathbb{C})/\text{GL}_{n-m}(\mathbb{C})) \rightarrow \text{Hol}_1^*(\mathbb{CP}^m, \mathbb{CP}^n).$$

Simple computation shows that $\beta \circ \gamma = \text{id}$, $\gamma \circ \beta = \text{id}$, and γ is a homeomorphism. Hence, $\text{Hol}_1^*(\mathbb{CP}^m, \mathbb{CP}^n) \cong \mathbb{C}^m \times (\text{GL}_n(\mathbb{C})/\text{GL}_{n-m}(\mathbb{C})) \simeq W_{n,m}$. \square

Corollary 2.4 ([10]). *There are homotopy equivalences*

$$\begin{cases} \text{Hol}_1^*(\mathbb{CP}^n, \mathbb{CP}^n) \simeq U_n, \\ \text{Hol}_1(\mathbb{CP}^n, \mathbb{CP}^n) \simeq P(SU_{n+1}) = SU_{n+1}/\Delta_{n+1}. \end{cases} \quad \square$$

3 The space $\text{Hol}(\mathbb{CP}^m, \mathbb{CP}^n)$.

Recently, J. Mostovoy [12] obtained the remarkable important result concerning the topology of $\text{Hol}(\mathbb{CP}^m, \mathbb{CP}^n)$. In this section we recall his result, and for this purpose, we study the restriction fibration sequence

$$F_d(m, n) \rightarrow \text{Map}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) \xrightarrow{r} \text{Map}_d^*(\mathbb{CP}^{m-1}, \mathbb{CP}^n),$$

where the map r is defined by the restriction $r(f) = f|_{\mathbb{CP}^{m-1}}$, $F_d(m, n)$ denotes the fiber of r defined by

$$F_d(m, n) = r^{-1}(\psi_d^{m-1, n}) = \{f \in \text{Map}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) : f|_{\mathbb{CP}^{m-1}} = \psi_d^{m-1, n}\},$$

and $\psi_d^{m, n} \in \text{Map}_d^*(\mathbb{CP}^m, \mathbb{CP}^n)$ is the holomorphic map defined by

$$\psi_d^{m, n}([x_0 : \dots : x_m]) = [(x_0)^d : \dots : (x_m)^d : 0 : \dots : 0]$$

for $[x_0 : \dots : x_m] \in \mathbb{CP}^m$.

We choose it as the basepoint of $\text{Map}_d^*(\mathbb{CP}^m, \mathbb{CP}^n)$. We remark that there is a homotopy equivalence $F_d(m, n) \simeq \Omega^{2m} \mathbb{CP}^n$ ([16]).

Let $H_d(m, n) \subset \text{Hol}_d^*(\mathbb{CP}^m, \mathbb{CP}^n)$ be the subspace defined by $H_d(m, n) = F_d(m, n) \cap \text{Hol}_d^*(\mathbb{CP}^m, \mathbb{CP}^n)$. We investigate the homotopy types of the subspaces $H_d(m, n)$, $\text{Hol}_d^*(\mathbb{CP}^m, \mathbb{CP}^n)$ and $\text{Hol}_d(\mathbb{CP}^m, \mathbb{CP}^n)$ with the corresponding inclusion maps

$$\begin{cases} i'_d : H_d(m, n) \rightarrow F_d(m, n), & i_d : \text{Hol}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \text{Map}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) \\ j_d : \text{Hol}_d(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \text{Map}_d(\mathbb{CP}^m, \mathbb{CP}^n). \end{cases}$$

Theorem 3.1 (J. Mostovoy, [13]). *If $2 \leq m \leq n$, the inclusion maps*

$$\begin{cases} i'_d : H_d(m, n) \rightarrow F_d(m, n), & i_d : \text{Hol}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \text{Map}_d^*(\mathbb{CP}^m, \mathbb{CP}^n) \\ j_d : \text{Hol}_d(\mathbb{CP}^m, \mathbb{CP}^n) \rightarrow \text{Map}_d(\mathbb{CP}^m, \mathbb{CP}^n) \end{cases}$$

are homotopy equivalences through dimension $D(d; m, n)$ when $m < n$, and homology equivalences through dimension $D(d; m, n)$ when $m = n$.

□

Since $\lim_{d \rightarrow \infty} D(d; m, n) = \infty$, we may regard $H_d(m, n)$ and $\text{Hol}(\mathbb{CP}^m, \mathbb{CP}^n)$ as finite dimensional homotopy (or homology) models for the infinite dimensional spaces $\Omega^{2m} \mathbb{CP}^n$ and $\text{Map}(\mathbb{CP}^m, \mathbb{CP}^n)$, respectively. We know

that the Atiyah-Jones type Theorem holds for several other cases, and the homotopy stability is usually satisfied for these cases (cf. [2], [3], [5], [6], [9], [19]). So one may expect that the homotopy stability may hold even if $m = n$. We consider the case $m = n$ in the next section.

4 The case $m = n$.

In this section we consider the space $\text{Hol}(\mathbb{CP}^n, \mathbb{CP}^n)$ when $m = n$. From now on, we write $F_d(n) = F_d(n, n)$ and $H_d(n) = H_d(n, n)$.

First, recall the following two results.

Theorem 4.1 ([21]). (1) *If $m = n$, there are isomorphisms*

$$\begin{cases} \pi_1(\text{Hol}_d^*(\mathbb{CP}^n, \mathbb{CP}^n)) = \pi_1(\text{Hol}_d^*) \cong \mathbb{Z} \\ \pi_1(\text{Hol}_d(\mathbb{CP}^n, \mathbb{CP}^n)) = \pi_1(\text{Hol}_d) \cong \mathbb{Z}/(n+1)d^n \end{cases}$$

(2) *The inclusion maps induce isomorphisms*

$$\begin{cases} i_{d*} : \pi_1(\text{Hol}_d^*) \xrightarrow{\cong} \pi_1(\text{Map}_d^*) = \pi_1(\text{Map}_d^*(\mathbb{CP}^n, \mathbb{CP}^n)), \\ j_{d*} : \pi_1(\text{Hol}_d) \xrightarrow{\cong} \pi_1(\text{Map}_d) = \pi_1(\text{Map}_d(\mathbb{CP}^n, \mathbb{CP}^n)). \end{cases} \quad \square$$

Theorem 4.2 ([22]). *If $n \geq 1$ and $d \geq 0$ be integers, the inclusion $i'_d : H_d(n) \rightarrow F_d(n) \simeq \Omega^{2n}\mathbb{CP}^n$ induces an isomorphism*

$$i'_{d*} : \pi_1(H_d(n)) \xrightarrow{\cong} \pi_1(F_d(n)) \cong \pi_1(\Omega^{2n}\mathbb{CP}^n) \cong \mathbb{Z}.$$

Sketch proof. If we use an easy diagram chasing and Theorem 4.1, we can show that $\pi_1(H_d(n)) \cong \mathbb{Z}$. Then the assertion follows from Theorem 3.1 and Hurewicz Theorem. \square

If we use the above two results and some group actions, we obtain the following two results.

Theorem 4.3 ([22]). *If $n \geq 2$, the inclusion $i'_d : H_d(n) \rightarrow F_d(n) \simeq \Omega^{2n}\mathbb{CP}^n$ is a homotopy equivalence through dimension $D(d, n)$, where we take $D(d, n) = D(d; n, n) = \lfloor \frac{d+1}{2} \rfloor$.* \square

Theorem 4.4 ([22]). *If $n \geq 2$, the inclusion maps*

$$\begin{cases} i_d : \text{Hol}_d^*(\mathbb{CP}^n, \mathbb{CP}^n) \rightarrow \text{Map}_d^*(\mathbb{CP}^n, \mathbb{CP}^n) \\ j_d : \text{Hol}_d(\mathbb{CP}^n, \mathbb{CP}^n) \rightarrow \text{Map}_d(\mathbb{CP}^n, \mathbb{CP}^n) \end{cases}$$

are homotopy equivalences through dimension $D(d, n) = \lfloor \frac{d+1}{2} \rfloor$. \square

Proof of Theorem 1.2. The assertion follows from Theorem 3.1 and Theorem 4.4. \square

5 The orbit spaces.

Finally, in this section, we consider the right $\text{PGL}_{n+1}(\mathbb{C})$ action on $\text{Hol}_d(\mathbb{CP}^n, \mathbb{CP}^n)$ given by the matrix multiplication

$$\begin{aligned} \text{Hol}_d(\mathbb{CP}^n, \mathbb{CP}^n) \times \text{PGL}_{n+1}(\mathbb{C}) &\longrightarrow \text{Hol}_d(\mathbb{CP}^n, \mathbb{CP}^n). \\ ([f_0 : \dots : f_n], A) &\longrightarrow [f_0 : \dots : f_n] \cdot A \end{aligned}$$

We denote by $X_d(n)$ the orbit space $X_d(n) = \text{Hol}_d(\mathbb{CP}^n, \mathbb{CP}^n) / \text{PGL}_{n+1}(\mathbb{C})$.

Theorem 5.1 (R. Milgram, [11]). *If $n = 1$, there is a homeomorphism $X_d(1) \cong P(\mathcal{F}_d)$, where \mathcal{F}_d denotes the space consisting of all $(d \times d)$ non-singular Toeplitz matrices.* \square

Then our main result of this section is as follows.

Theorem 5.2 ([21]). (1) $\pi_1(X_d(n)) \cong \mathbb{Z}/d^n$.

(2) *There is a fibration sequence (up to homotopy)*

$$(*) \quad SU_{n+1} \rightarrow \widetilde{X_d(n)} \rightarrow \widetilde{\text{Hol}_d(\mathbb{CP}^n, \mathbb{CP}^n)},$$

where \widetilde{Y} denotes the universal covering of a space Y .

(3) *If $n = 1$, the fibration $(*)$ is trivial and there is a homotopy equivalence $\widetilde{X_d(1)} \simeq S^3 \times \widetilde{\text{Hol}_d(\mathbb{CP}^1, \mathbb{CP}^1)}$.* \square

Corollary 5.3 (J. Havlicek, [8]; [21]). $\pi_1(X_d(1)) \cong \mathbb{Z}/d$. \square

Acknowledgements. The author is indebted to Professors M. A. Guest, A. Kozłowski and J. Mostovoy for numerous helpful conversations concerning the topology of configuration spaces and valuable suggestions. The author is partially supported by a grant from the Ministry of Education.

References

- [1] M. F. Atiyah and J. D. S. Jones, Topological aspects of Yang-Mills theory, *Commun. Math. Phys.* **59** (1978), 97–118.
- [2] C. P. Boyer, J. C. Hurtubise, B. M. Mann and R. J. Milgram, The topology of the space of rational maps into generalized flag manifolds, *Acta Math.* **173** (1994), 61–101.
- [3] M. A. Guest, Topology of the space of rational curves on a toric variety, *Acta Math.* **174** (1995), 119–145.
- [4] M. A. Guest, A. Kozłowski, M. Murayama, and K. Yamaguchi, The homotopy type of spaces of rational functions, *J. Math. Kyoto Univ.* **35** (1995), 631–638.
- [5] M. A. Guest, A. Kozłowski, and K. Yamaguchi, The topology of spaces of coprime polynomials, *Math. Z.* **217** (1994), 435–446.
- [6] M. A. Guest, A. Kozłowski and K. Yamaguchi, Spaces of polynomials with roots of bounded multiplicity, *Fund. Math.* **116** (1999), 93–117.
- [7] J. Harris, *Algebraic Geometry*, Springer-Verlag, 1993.
- [8] J. W. Havlicek, The cohomology of holomorphic self maps of the Riemann sphere, *Math. Z.* **218** (1995), 179–190.
- [9] A. Kozłowski and K. Yamaguchi, Topology of complements of discriminants and resultants, *J. Math. Soc. Japan* **52** (2000), 949–959.
- [10] A. Kozłowski and K. Yamaguchi, Spaces of holomorphic maps between complex projective spaces of degree one, *Topology Appl.* **132** (2003), 139–145.
- [11] R. Milgram, The structure of spaces of Toeplitz matrices, *Topology* **36** (1997), 1155–1192.
- [12] J. Mostovoy, Spaces of rational loops on a real projective space, *Trans. Amer. Math. Soc.* **353** (2001), 1959–1970.

- [13] J. Mostovoy, Spaces of rational maps and the Stone-Weierstrass Theorem, preprint, arXiv:math. AT/0307103, (to appear in Topology).
- [14] J. M. Møller, On spaces of maps between complex projective spaces, Proc. Amer. Math. Soc. **91** (1984), 471–476.
- [15] Y. Ono and K. Yamaguchi, Group actions on spaces of rational functions, Publ. Res. Inst. Math. Sci. **39** (2003), 173–181.
- [16] S. Sasao, The homotopy of $\text{Map}(\mathbb{CP}^m, \mathbb{CP}^n)$, J. London Math. Soc. **8** (1974), 193–197.
- [17] G. B. Segal, The topology of spaces of rational functions, Acta Math. **143** (1979), 39–72.
- [18] V. A. Vassiliev, Complements of discriminants of smooth maps, topology and applications, Amer. Math. Soc., Translations of Math. Monographs **98**, 1992 (revised edition 1994).
- [19] K. Yamaguchi, Spaces of holomorphic maps with bounded multiplicity, Quart. J. Math. **52** (2001), 249–259.
- [20] K. Yamaguchi, Configuration space models for spaces of maps from a Riemann surface to complex projective space, Publ. Res. Inst. Math. Sci. **39** (2003), 535–543.
- [21] K. Yamaguchi, Fundamental groups of spaces of holomorphic maps and group actions, J. Math. Kyoto Univ. **44** (2004), 479–492.
- [22] K. Yamaguchi, Spaces of holomorphic self-maps of the complex projective spaces, preprint.